

AUGTYRO® (repotrectinib) FDA-APPROVED TO TARGET MY **NTRK+ CANCER**



Actor portrayal

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).

What is AUGTYRO?

AUGTYRO (repotrectinib) is an oral prescription medicine used to treat adults and children 12 years and older with **solid tumors (cancer)** that:

- ✓ Are caused by certain abnormal *NTRK* genes, **AND**
- ✓ Have spread to other parts of the body or if surgery to remove your cancer is likely to cause severe complications, **AND**
- ✓ Have grown or spread after any other treatment or there is no satisfactory alternative treatment option.

It is not known if AUGTYRO is safe and effective in children younger than 12 years of age with *NTRK*-positive solid tumors.




Talk to your healthcare team for more information about what it means to have an abnormal *NTRK* genes.

SUMMARY OF WARNINGS AND PRECAUTIONS


AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.





Many tumors in different areas of the body can be *NTRK*+

This chart does not include all types of possible *NTRK*+ cancers in adults and children and not all cancer types were studied in the AUGTYRO clinical trial.



To see a list of cancer types that were included in the study, please refer to the [U.S. Full Prescribing Information](#).

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).

 **AUGTYRO™**
(repotrectinib) 40 mg capsules

ARE YOU READY FOR THIS?

A cancer diagnosis can undoubtedly catch you off guard, especially a rare type of cancer known as *NTRK*+ cancer, or *NTRK* gene fusion-positive cancer. It may bring about uncertainty and fill your head with questions. The following pages offer guidance for a next step, which may involve a treatment targeted to help fight *NTRK*+ cancer in pill form—AUGTYRO, a treatment that may help tumors shrink or disappear (see page 5 for more information).


Look to this guide and your healthcare team for answers. You may start at the beginning and learn about an oral, chemo-free treatment, targeted to treat *NTRK*+ cancer. Or, jump to other sections about side effects, dosing, and support. **It's important to learn how AUGTYRO works differently than chemotherapy by targeting your cancer. It's a treatment designed to find certain cancer cells and help turn off their growth signals.**

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).





Less than **1%** of cancers are *NTRK+*

A TARGETED THERAPY FOR *NTRK+* CANCER

Whether you're considering targeted therapy for the first time or you've had it before, AUGTYRO offers people like yourself a specific treatment option.

AUGTYRO was studied in people whose cancer was *NTRK+*. This study included people in two important groups: 40 people who had no previous treatment with targeted therapy, and 48 people who were previously treated with targeted therapy.



AUGTYRO has been shown to provide results for people with *NTRK+* cancer.



AUGTYRO, taken orally in pill form, is targeted to treat *NTRK+* cancer. Response to treatment may vary.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).





Study results

in people with *NTRK*+ cancer who were given AUGTYRO

NTRK+ tumors can occur in many different types of cancer throughout the body. The AUGTYRO study included people with non-small cell lung cancer, salivary gland cancer, thyroid cancer, soft-tissue sarcoma, breast cancer, glioblastoma, and several other cancers.

	Previously untreated with targeted therapy (of 40 people)	Previously treated with targeted therapy (of 48 people)
Tumors shrank or disappeared* (Overall response)	58%	50%
Tumors shrank (Partial response)	43%	50%
Tumors disappeared* (Complete response)	15%	0%

STILL RESPONDING AT 12 MONTHS

83% of previously
untreated people

42% of previously
treated people

Individual experience may vary. If tumors disappear during treatment, continuing treatment is still necessary.

*The disappearance of many measurable tumors in response to treatments does not necessarily mean the cancer has been cured.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



WHAT YOU KNOW **MATTERS.**

WHAT YOU DO NEXT
MATTERS EVEN MORE.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



Actor portrayal

 **AUGTYRO™**
(repotrectinib) 40 mg capsules

NOT ALL **CANCERS ARE THE SAME.**

To determine your specific type, your cancer may have been tested for a unique marker, called a biomarker. Some cancers, like yours, test positive for *NTRK*. While your diagnosis is a rare type of cancer, knowing your biomarker helped your doctor determine that AUGTYRO may be right for you.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



BIOMARKERS MAY HAVE THE DETAILS



Biomarkers (also called tumor markers) can offer valuable insight into your cancer. They are determined through biomarker testing. This is when your doctor checks a blood or tissue sample for certain genes, proteins, or other markers that could be a sign of a specific disease or cancer.

Here's why knowing your biomarker is important:

- Biomarkers give you and your doctor details about your cancer so that your doctor can accurately diagnose it.
- They help determine the best plan of action. Think of biomarkers as keys that help you and your doctor unlock treatment options.



All in all, biomarker testing (also referred to as tumor testing or molecular testing) helped you and your doctor determine that you have *NTRK+* cancer and that **AUGTYRO may be an effective treatment for you.**

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



WHAT TYPE OF TREATMENT IS AUGTYRO[®] (repotrectinib)?

AUGTYRO is a targeted therapy to treat *NTRK*+ cancer. It's a prescription oral treatment that's designed to help identify certain cancer cells, and shut off their "grow" signals.

AUGTYRO is not chemotherapy—
it's a targeted *NTRK*+ treatment.



AUGTYRO is a targeted therapy which means it zeroes in on proteins related to the *NTRK* gene. Most types of targeted therapies like AUGTYRO help fight specific proteins that cause tumors to grow and spread throughout the body.

AUGTYRO is a precise way of helping fight *NTRK*+ cancer.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



TAKING AUGTYRO[®] (repotrectinib)



For the first 14 days, you will start with a dose of 1 pill a day until your doctor recommends otherwise.



On Day 15, you will start taking 1 pill in the morning and 1 pill in the evening every day until your doctor recommends otherwise.

A 40-mg dosing option is also available.

Take AUGTYRO as directed

If you miss a dose or vomit at any time after taking a dose of AUGTYRO, do not take an extra dose. Take your next dose at the regularly scheduled time. Do not take 2 doses at the same time to make up a missed or vomited dose.

	AUGTYRO treats <i>NTRK</i> + cancer with an oral treatment in pill form that you can take from the comfort of your home.
	AUGTYRO may be taken with or without food. It should be swallowed whole with water at about the same time each day as prescribed. Avoid eating grapefruit or drinking grapefruit juice during your treatment with AUGTYRO. Remember not to open, crush, chew, or dissolve the capsule. Do not take a capsule if it is broken, cracked, or damaged.
	From Day 1 to Day 14, your doctor will start you on a once-daily dose. Starting on Day 15, the dose may be increased to twice daily.
	If you are experiencing any side effects, see page 12 for details and contact your care team. Your healthcare team will work closely with you to help manage these side effects with changes in dosing. In some cases they may temporarily or permanently stop treatment.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).





STICKING TO YOUR SCHEDULE

It's necessary to take AUGTYRO as prescribed by your doctor. This will help enable your treatment to be effective as you fight *NTRK+* cancer. Below are ways to help you remember to take a dose:



Develop a routine

Take AUGTYRO at about the same time each day to make it easier to remember. For example, at 8 AM and 8 PM. Or maybe before you brush your teeth in the morning and before you brush at night.



Give yourself a reminder

Place a note where you're likely to see it or set an alarm on your phone.



Use a child-resistant pill box

Keep track of your dosing schedule with a pill box labeled with the days of the week.



Record each dose

Use a calendar to check off when you've taken AUGTYRO each day.



Be prepared when traveling

Carry your AUGTYRO with you.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



POSSIBLE SIDE EFFECTS TO LOOK OUT FOR

Now that you've learned how AUGTYRO works, adjusting to your new "normal" may involve side effects. It's important to be mindful, track changes, and assess how you're feeling.

Rely on your care team

To have helpful conversations with them, remember to let your team know if you are experiencing any new or different symptoms at any time during your treatment.

AUGTYRO may cause serious side effects, including:

- Central nervous system (CNS) effects
- Lung problems (pneumonitis)
- Liver problems
- Muscle problems
- Increased uric acid in your blood (hyperuricemia)
- Bone fractures

The most common side effects of AUGTYRO include:

- Dizziness
- Change in sense of taste
- Feeling of numbness or tingling in your arms or legs
- Constipation
- Shortness of breath
- Tiredness
- Trouble with balance, coordination, and walking
- Problems with thinking, such as forgetfulness or confusion, memory problems, and hallucinations
- Muscle weakness
- Nausea

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



The most common side effect? Dizziness.

If at any point in your treatment you experience dizziness, let your healthcare team know. Do not drive or operate machinery until you know how AUGTYRO affects you. If you experience dizziness, do not drive or operate machinery until your symptoms have resolved.

Here are some things you can do that may help manage dizziness:

- ☒ Drink plenty of fluids daily. Aim for 8 to 12 eight-ounce glasses of water or other fluids every day.
- ☒ Avoid drinking a lot of caffeinated beverages including coffee, tea, and soda.
- ☒ Avoid moving suddenly. For example, sit up carefully from a lying position. If you get dizzy when you stand up, hold a chair or table for balance and stand up more slowly than usual.
- ☒ Sit or lie down immediately. The possibility of losing your balance could lead to falling or serious injury.

The tips provided here are for informational purposes only and are not meant to replace a physician's medical advice. Please contact your healthcare team if you are experiencing dizziness and ask your doctor if these tips may be right for you.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



HELP YOU STAY YOU

During this journey, remember that you're not alone. Reach out to your healthcare team, caregivers, friends, and loved ones. Here are some organizations that offer support to help you stay you.*

- American Cancer Society
cancer.org
- CancerCare
cancercare.org
- Cancer Support Community
cancersupportcommunity.org
- NTRKers
ntrkers.org

*Bristol Myers Squibb does not endorse these organizations. The information/links provided by Bristol Myers Squibb are meant for informational purposes only and are not meant to replace a physician's medical advice.



Actor portrayals

Feel seen, heard, supported. **Always.**

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).

 **AUGTYRO™**
(repotrectinib) 40 mg capsules

WHAT TO DISCUSS WITH YOUR DOCTOR

Understandably, you may still have some questions about *NTRK*+ cancer, AUGTYRO, or both. Here are some questions you can discuss with your doctor.

- How does AUGTYRO work?
- Do you think AUGTYRO will help me?
- What are the potential side effects that I may experience?
- How often will I need scans or follow-up visits during treatment?
- How will I know if AUGTYRO is working?
- Will this treatment impact my daily life in any way?

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).





Actor portrayals

HOW TO GET AUGTYRO

You can get AUGTYRO through certain specialty pharmacies that handle treatments, including AUGTYRO, which are not stocked at regular neighborhood pharmacies.

Most specialty pharmacies deliver medications by mail.

Here are some steps your doctor's office and the specialty pharmacy must do before AUGTYRO can be shipped to you:

- Make sure that your health insurance plan has approved AUGTYRO.
- Discuss shipping details with you.
- Collect your co-pay.



Your specialty pharmacy will call to confirm the above details. Make sure you save their number in your phone so you can answer right away or return their calls. This will help you get your AUGTYRO treatment on time.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



At Bristol Myers Squibb, We Provide Support With Purpose

You are the reason behind what we do. The BMS Access Support® program is focused on helping you access your prescribed BMS medicine. If you were prescribed AUGTYRO, you are eligible to enroll in BMS Access Support. The program can help you with:



Coverage Assistance

Upon your request, BMS Access Support may be able to help you review your insurance benefits and understand your coverage for AUGTYRO.*

If your insurer requires prior authorization, BMS Access Support may be able to provide you and your doctor information about this requirement.

In the event of a coverage delay or denial, you may be eligible for a bridge program to cover your treatment.† Please call BMS Access Support at 1-800-861-0048 for more information about offerings for your medication.



Financial Support

If you are an eligible commercially-insured patient, you may pay as little as \$0 per one-month supply.†

If you are insured through a government program or you do not have insurance, BMS Access Support can provide you with information regarding independent charitable foundations.‡



Free Trial Offer

A Free Trial Offer may be available if you were newly prescribed AUGTYRO.†



Educational Resources

Understanding the insurance process and accessing your treatment can be challenging. BMS Access Support offers a library of resources that can help you better understand your healthcare coverage and guide you through steps in your treatment journey.

*The accurate completion and submission of reimbursement- or coverage-related documentation to the patient's insurance plan is the responsibility of the provider and patient. Bristol Myers Squibb and its agents cannot guarantee coverage for any medication or treatment.

†Restrictions apply. Please [click here](#) for full Terms and Conditions, including complete eligibility requirements.

‡It is important to note that charitable foundations are independent from Bristol-Myers Squibb Company and have their own eligibility criteria and evaluation process. Bristol Myers Squibb cannot guarantee that a patient will receive assistance.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).

We're here for you.
Coverage assistance, educational resources, and financial support options may be available through **BMS Access Support**



Call a Patient Access Specialist at
1-800-861-0048, 8 AM to 8 PM ET, Monday - Friday



Find resources and enrollment information at
www.BMSAccessSupport.com

 **AUGTYRO™**
(repotrectinib) 40 mg capsules

**Important Facts
about AUGTYRO™
(repotrectinib)**

This is a summary of important information that you need to know about AUGTYRO. Your healthcare team can work with you to help answer any questions you may have about this medication. [Keep this document in a safe place, so you can refer to it before and during your treatment.](#)

Look out for the following icons as you read:

 Talk to your healthcare team

 Call a healthcare provider right away


 Helpful information to remember

What is AUGTYRO?

AUGTYRO (repotrectinib) is an oral prescription medicine used to treat adults and children 12 years and older with **solid tumors (cancer)** that:

- ✓ Are caused by certain abnormal *NTRK* genes, **AND**
- ✓ Have spread to other parts of the body or if surgery to remove your cancer is likely to cause severe complications, **AND**
- ✓ Have grown or spread after any other treatment or there is no satisfactory alternative treatment option.

It is not known if AUGTYRO is safe and effective in children younger than 12 years of age with *NTRK*-positive solid tumors.


 **Talk to your healthcare team** for more information about what it means to have an abnormal *NTRK* genes.

What is the most important information I should know about AUGTYRO?

AUGTYRO may cause serious side effects. A serious side effect is a side effect that can sometimes become life-threatening and can lead to death. The serious side effects of AUGTYRO may include:


Central nervous system (CNS) problems. Symptoms to look out for include:

- dizziness
- vertigo
- changes in mood, such as anxiety, irritability, and depression
- balance or coordination problems
- problems with thinking, such as forgetfulness or confusion
- seeing or hearing things that are not real (hallucinations)
- problems with concentration, attention, memory, and sleep

 **Call your healthcare provider right away** if you have any of these symptoms, or any worsening of CNS effects during treatment.

Lung problems (pneumonitis). AUGTYRO may cause pneumonitis (inflammation of the lungs), which may be severe or serious. New or worsening symptoms to look out for include:

- Dry cough (without mucus)
- Productive cough (with mucus)
- Wheezing
- Trouble breathing

 **Call your healthcare provider right away** if you have any of these symptoms, or any worsening of your current lung problems.

Liver problems. Your healthcare provider will do blood tests to check your liver function before starting treatment with AUGTYRO™, every 2 weeks for the first month, and as needed during treatment. Symptoms of liver problems include:


- your skin or the white part of your eyes turns yellow
- dark or “tea-colored” urine
- light-colored stools (bowel movements)
- loss of appetite
- nausea or vomiting
- pain on the upper right side of your stomach area

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



 **Call your healthcare provider right away** if you develop any of these symptoms of liver problems.

Muscle problems. Your healthcare provider will do blood tests before starting treatment with AUGTYRO, every 2 weeks for the first month, and as needed during treatment. Symptoms of muscle problems include:


- unexplained muscle pain or muscle pain that does not go away
- tenderness
- weakness

 **Call your healthcare provider right away** if you get new or worsening signs and symptoms of muscle problems.

Increased uric acid level in your blood (hyperuricemia). AUGTYRO may cause an excess of uric acid in your blood. Your healthcare provider will do tests before and during your treatment with AUGTYRO to check the uric acid level in your blood. Your healthcare provider may prescribe medications if you have high blood uric acid levels. Tell your healthcare provider if you experience symptoms of increased uric acid, including:

- red, hot, tender, or swollen joints, especially in your big toe
- decrease in your amount of urine or no urine at all
- nausea or vomiting
- pain in your stomach-area or sides
- pink or brown urine or blood in your urine

Bone fractures. AUGTYRO may increase your risk for bone fractures. Bone fractures may happen with or without a fall or other injury.

 **Call your healthcare provider right away** if you have pain, changes in movement, or bone abnormalities.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).

What should I discuss with my healthcare team before starting treatment?

 **Talk to your healthcare provider about all your medical conditions,** including if you:

- have nervous system (neurological) problems
- have lung or breathing problems other than lung cancer
- have liver or kidney problems

 **Talk to your healthcare team about all the medicines you are taking,** including:

- prescription medicines
- over-the-counter medicines
- vitamins
- herbal supplements

Taking AUGTYRO with certain other medicines may:

- affect the amount of AUGTYRO or other medicines in your blood,
- cause side effects, and/or
- affect the way that AUGTYRO or other medicines work.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

These are not all the topics you should discuss with your healthcare team. Discuss any questions you may have with your healthcare team.

What should I discuss with my healthcare team about pregnancy, birth control, and breastfeeding before starting treatment?

 **Talk to your healthcare team** if:



You are pregnant or plan to become pregnant – AUGTYRO™ may harm your unborn baby


 **AUGTYRO™**
(repotrectinib) 40 mg capsules

If you are able to become pregnant, your healthcare team should do a pregnancy test before you start treatment with AUGTYRO.

If you are able to become pregnant, you should use effective non-hormonal birth control (contraception) during treatment and for 2 months after your last dose of AUGTYRO.


Birth control methods that contain hormones (such as birth control pills, injections, or transdermal system patches) may not work as well during treatment with AUGTYRO.

 **Talk to your healthcare team** about **birth control methods** that may be right for you.

 **Call your healthcare provider right away** if you become pregnant or think you may be pregnant during treatment with AUGTYRO.

You are a male with a female partner who is able to become pregnant

You should use effective birth control during treatment with AUGTYRO and for 4 months after your last dose.

 **Call your healthcare provider right away** if your partner becomes pregnant during treatment with AUGTYRO.



You are breastfeeding or plan to breastfeed

Do not breastfeed during treatment and for 10 days after the last dose of AUGTYRO. It is not known if AUGTYRO passes into your breast milk. Talk to your healthcare team about the best way to feed your baby during this time.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).

What are the most common side effects of AUGTYRO?


The most common side effects of AUGTYRO include:


- Dizziness
- Change in sense of taste
- Feeling of numbness or tingling in your arms or legs
- Constipation
- Shortness of breath
- Tiredness
- Trouble with balance, coordination, and walking
- Problems with thinking, such as forgetfulness or confusion, memory problems, and hallucinations
- Muscle weakness
- Nausea

These are not all the possible side effects. **Talk to your doctor** for more information about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA by visiting www.fda.gov/medwatch or calling 1-800-FDA-1088.



How should AUGTYRO™ be taken?

 **Take AUGTYRO exactly as your healthcare provider tells you.** The number of pills that you take will depend on your dose. Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with AUGTYRO if you develop side effects. Do not change your dose or stop taking AUGTYRO without first talking to your healthcare provider.



1.8 cm


AUGTYRO is a capsule with a white opaque cap and body, filled with a white to off-white powder. "REP 40" is printed in blue text on the cap.

Inactive ingredients: microcrystalline cellulose, sodium lauryl sulfate, croscarmellose sodium, and colloidal silicon dioxide.

Capsule coating inactive ingredients: gelatin and titanium dioxide.

Printing ink: shellac and FD & C blue #2 aluminum lake.

OR



1.8 cm

AUGTYRO is a capsule with a blue opaque cap and body, filled with a white to off-white powder. "REP 160" is printed in white text on the cap.

Inactive ingredients: microcrystalline cellulose, sodium lauryl sulfate, croscarmellose sodium, magnesium stearate, and colloidal silicon dioxide.

Capsule coating inactive ingredients: gelatin and titanium dioxide.

Printing ink: shellac, titanium dioxide, and FD & C blue #1.

- ✓ **Do** take AUGTYRO with or without food
- ✓ **Do** take AUGTYRO at about the same time each day, such as in the morning or in the evening*
- ✓ **Do** swallow AUGTYRO capsules **whole** with water

**Setting an alarm clock or a calendar notification is a great way to remember to take medication.*


- ✗ **Do not** open, crush, chew, or dissolve the capsule. **Do not** take broken, cracked, or damaged capsules
- ✗ **Do not** eat grapefruit or drink grapefruit juice during treatment with AUGTYRO
- ✗ **Do not** drive or operate machinery until you know how AUGTYRO affects you and you no longer have symptoms

What if I miss a dose of AUGTYRO?

If you miss a dose or vomit at any time after taking a dose of AUGTYRO, **do not take an extra dose.** Just skip the dose and take your next dose at the regularly scheduled time. Do not take 2 doses at the same time to make up a missed or vomited dose.

How do I store AUGTYRO?


AUGTYRO capsules should be stored at room temperature between 68°F to 77°F (20°C to 25°C). **Keep AUGTYRO and all medicines out of the reach of children.**

 For more information, please see accompanying U.S. Full Prescribing Information and Patient Information for AUGTYRO. Talk to your healthcare team for more information about this medication.

SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).



21



Actor portrayal



We're here throughout your treatment journey.
Scan here to learn more about AUGTYRO
and sign up for AUGTYRO™ *with You* to receive support,
tools, and educational resources.



Call us toll-free at **1-855-436-5866**



© 2024 Turning Point Therapeutics, Inc., a Bristol Myers Squibb company.
AUGTYRO®, AUGTYRO® *with You*, and the related logos are trademarks of Turning Point
Therapeutics, Inc. Access Support® is a registered trademark of Bristol-Myers Squibb
Company. All other trademarks are property of their respective owners.
3600-US-2400279 10/24

**SUMMARY OF WARNINGS AND
PRECAUTIONS**

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

Please see Important Safety Information on pages 18-21. Click on links for U.S. Full Prescribing Information and Patient Information.

