

AUGTYRO™ (repotrectinib)  
IS MADE SPECIFICALLY FOR MY

**ROS** +

AUGTYRO is designed  
for your type of lung cancer.

Actor portrayal

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**

### What is AUGTYRO?

AUGTYRO (repotrectinib) is an oral prescription medicine used to treat **non-small cell lung cancer (NSCLC) in adults** that:

- ✓ Has spread within your chest or to other parts of the body, **AND**
- ✓ Is caused by an **abnormal ROS1 gene**

It is not known if AUGTYRO is safe and effective in children.



**Talk to your healthcare team** for more information about what it means to have an abnormal ROS1 gene.

### SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

 **AUGTYRO™**  
(repotrectinib) 40 mg capsules





People with *ROS1+* lung cancer tend to be younger (about age 50) than lung cancer patients in general (age 70) and have little to no smoking history.

## **SUMMARY OF WARNINGS AND PRECAUTIONS**

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**



**AUGTYRO™**  
(repotrectinib) 40 mg capsules

Actor portrayal

## ARE YOU READY FOR THIS?

A lung cancer diagnosis may undoubtedly catch you off guard, especially *ROS1*+ non-small cell lung cancer, a rare form of lung cancer. It may bring about uncertainty and fill your head with questions. The following pages offer guidance for a next step, which may involve a treatment tailored to help fight *ROS1*+ lung cancer in pill form—AUGTYRO, a treatment that may help lung tumors shrink or disappear (see page 5 for more information).

Look to this guide and your healthcare team for answers. You may start at the beginning and learn about an oral, chemo-free treatment, tailored for *ROS1*+ lung cancer. Or, jump to other sections about side effects, dosing, and support. **It's important to learn how AUGTYRO works differently than chemotherapy by targeting your cancer. It's a treatment designed to find certain cancer cells and help turn off their growth signals.**

### SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**





1%-2% of adults with non-small cell lung cancer are diagnosed with *ROS1*+ lung cancer.

## START WITH A TREATMENT TARGETED FOR *ROS1*

People with *ROS1*+ non-small cell lung cancer, like yourself, can be treated with AUGTYRO as their first targeted therapy.

In a study, 71 people with *ROS1*+ lung cancer were given AUGTYRO as their first targeted therapy. They received AUGTYRO starting at 160 mg orally once daily for 14 days, and then their dose was increased to 160 mg twice daily until their doctor recommended otherwise.



AUGTYRO has been shown to provide results for people with *ROS1*+ lung cancer.



AUGTYRO, taken orally in pill form, is tailored to treat *ROS1*+ lung cancer. Response to treatment may vary.

### SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**

 **AUGTYRO™**  
(repotrectinib) 40 mg capsules





Actor portrayals

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**

## Study Results

in people given AUGTYRO as their first targeted therapy

### Nearly **3 YEARS**

Half of the people taking AUGTYRO were still responding to treatment at 34 months.

### Less than **2 MONTHS**

For some, AUGTYRO started showing positive effects in people in a range of 1.4+, 42.4+ months.

Of 71 people with <i>ROS1+</i> NSCLC who participated in the study	
Tumors disappeared* (Complete response)	<b>6%</b>
Tumors shrank (Partial response)	<b>73%</b>
Tumors shrank or disappeared* (Overall response)	<b>79%</b>

Individual experience may vary. Some may notice improvements sooner or later than the average time frame. If tumors disappear during treatment, continuing treatment is still necessary.

\*The disappearance of many measurable tumors in response to treatments does not necessarily mean the cancer has been cured.



WHAT YOU KNOW **MATTERS.**

WHAT YOU DO NEXT  
**MATTERS EVEN MORE.**



## **SUMMARY OF WARNINGS AND PRECAUTIONS**

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**

 **AUGTYRO™**  
(repotrectinib) 40 mg capsules



## **NOT ALL LUNG CANCERS ARE THE SAME.**

To determine your specific type, your cancer may have been tested for a unique marker, called a biomarker. Some non-small cell lung cancers, like yours, test positive for *ROS1*.\* While your diagnosis is a rare type of lung cancer, knowing your biomarker helped your doctor determine that AUGTYRO may be right for you.

\*An error in the *ROS1* gene is one biomarker that doctors look for in non-small cell lung cancer.

## **SUMMARY OF WARNINGS AND PRECAUTIONS**

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**



# BIOMARKERS MAY HAVE THE DETAILS



Biomarkers (also called tumor markers) can offer valuable insight into your cancer. They are determined through biomarker testing. This is when your doctor checks a blood or tissue sample for certain genes, proteins, or other markers that could be a sign of a specific disease or cancer.

## Here's why knowing your biomarker is important:

- Biomarkers give you and your doctor details about your cancer so that your doctor can accurately diagnose it.
- They help determine the best plan of action. Think of biomarkers as keys that help you and your doctor unlock treatment options.



All in all, biomarker testing (also referred to as tumor testing or molecular testing) helped you and your doctor determine that you have *ROS1*+ lung cancer and that **AUGTYRO** may be an effective treatment for you.

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**





# WHAT TYPE OF TREATMENT IS AUGTYRO™ (repotrectinib)?

AUGTYRO is a targeted therapy tailored specifically to treat *ROS1*+ lung cancer. It's a prescription oral treatment that's designed to help identify certain cancer cells, and shut off their "grow" signals.

**AUGTYRO is not chemotherapy—**  
**it's a targeted *ROS1* treatment.**



AUGTYRO is a targeted therapy which means it zeroes in on *ROS1* proteins. Most types of targeted therapies like AUGTYRO help fight specific proteins that cause tumors to grow and spread throughout the body.

AUGTYRO is a precise way of helping fight *ROS1*+ lung cancer.

## SUMMARY OF WARNINGS AND PRECAUTIONS


AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**



# TAKING AUGTYRO™ (repotrectinib)

## Starting treatment

 **x4** Days 1-14 Your daily dose:  
**4 total pills a day.**

For the first 14 days, you will start with a dose of 4 pills once a day until your doctor recommends otherwise.

## Continuing treatment

 **x4**  **AM** +  **PM** Day 15 and throughout your treatment  
Your daily dose: **8 total pills a day.**

On Day 15, you will start taking 4 pills in the morning and 4 pills in the evening every day until your doctor recommends otherwise.

## Take AUGTYRO as directed

If you miss a dose or vomit at any time after taking a dose of AUGTYRO, do not take an extra dose. Take your next dose at the regularly scheduled time. Do not take 2 doses at the same time to make up a missed or vomited dose.



AUGTYRO treats *ROS1*+ lung cancer with an oral treatment in pill form that you can take from the comfort of your home.



AUGTYRO may be taken with or without food. It should be swallowed whole with water at about the same time each day as prescribed. Avoid eating grapefruit or drinking grapefruit juice during your treatment with AUGTYRO. Remember not to open, crush, chew, or dissolve the capsule. Do not take a capsule if it is broken, cracked, or damaged.



From Day 1 to Day 14, your doctor will start you on a once-daily dose. Starting on Day 15, the dose may be increased to twice daily.



If you are experiencing any side effects, see page 12 for details and contact your care team. Your healthcare team will work closely with you to help manage these side effects with changes in dosing. In some cases they may temporarily or permanently stop treatment.

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**

 **AUGTYRO™**  
(repotrectinib) 40 mg capsules



## STICKING TO YOUR SCHEDULE

It's necessary to take AUGTYRO as prescribed by your doctor. This will help enable your treatment to be effective as you fight *ROS1+* lung cancer. Below are ways to help you remember to take a dose:



### Develop a routine

Take AUGTYRO at about the same time each day to make it easier to remember. For example, at 8 AM and 8 PM. Or maybe before you brush your teeth in the morning and before you brush at night.



### Give yourself a reminder

Place a note where you're likely to see it or set an alarm on your phone.



### Use a child-resistant pill box

Keep track of your dosing schedule with a pill box labeled with the days of the week.



### Record each dose

Use a calendar to check off when you've taken AUGTYRO each day.



### Be prepared when traveling

Carry your AUGTYRO with you.

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**





# POSSIBLE SIDE EFFECTS TO LOOK OUT FOR

Now that you've learned how AUGTYRO works, adjusting to your new "normal" may involve side effects. It's important to be mindful, track changes, and assess how you're feeling.

## Rely on your care team

To have helpful conversations with them, remember to let your team know if you are experiencing any new or different symptoms at any time during your treatment.

AUGTYRO may cause serious side effects, including:

- Central nervous system (CNS) effects
- Lung problems (pneumonitis)
- Liver problems
- Muscle problems
- Increased uric acid in your blood (hyperuricemia)
- Bone fractures

The most common side effects of AUGTYRO include:

- Dizziness
- Change in sense of taste
- Feeling of numbness or tingling in your arms or legs
- Constipation
- Trouble with balance, coordination, and walking
- Shortness of breath
- Problems with thinking, such as forgetfulness or confusion, memory problems and hallucinations
- Tiredness
- Nausea

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**



## The most common side effect? Dizziness.

If at any point in your treatment you experience dizziness, let your healthcare team know. Do not drive or operate machinery until you know how AUGTYRO affects you. If you experience dizziness, do not drive or operate machinery until your symptoms have resolved.

## Here are some things you can do that may help manage dizziness:

- Drink plenty of fluids daily. Aim for 8 to 12 eight-ounce glasses of water or other fluids every day.
- Avoid drinking a lot of caffeinated beverages including coffee, tea, and soda.
- Avoid moving suddenly. For example, sit up carefully from a lying position. If you get dizzy when you stand up, hold a chair or table for balance and stand up more slowly than usual.
- Sit or lie down immediately. The possibility of losing your balance could lead to falling or serious injury.

The tips provided here are for informational purposes only and are not meant to replace a physician's medical advice. Please contact your healthcare team if you are experiencing dizziness and ask your doctor if these tips may be right for you.

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**



# HELP YOU STAY YOU

During this journey, remember that you're not alone. Reach out to your healthcare team, caregivers, friends, and loved ones. Here are some organizations that offer support to help you stay you.\*

- ROS1ders  
[theros1ders.org](http://theros1ders.org)
- LUNGeivity  
[lungevity.org](http://lungevity.org)
- GO2 for Lung Cancer  
[go2.org](http://go2.org)
- Lung Cancer Foundation of America  
[lcfamerica.org](http://lcfamerica.org)
- Lung Cancer Research Foundation  
[lungcancerresearchfoundation.org](http://lungcancerresearchfoundation.org)
- American Lung Association  
[lung.org](http://lung.org)
- A Breath of Hope Lung Foundation  
[youandlungcancer.com](http://youandlungcancer.com)

\*Bristol Myers Squibb does not endorse these organizations. The information/links provided by Bristol Myers Squibb are meant for informational purposes only and are not meant to replace a physician's medical advice.



Actor portrayals

Feel seen, heard, supported. **Always.**

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for U.S. Full Prescribing Information and Patient Information.**

 **AUGTYRO™**  
(repotrectinib) 40 mg capsules



# WHAT TO DISCUSS WITH YOUR DOCTOR

Understandably, you may still have some questions about *ROS1+* lung cancer, AUGTYRO, or both. Here are some questions you can discuss with your doctor. There is also room on this page for any additional notes you want to remember later.

- How does AUGTYRO work?
- Do you think AUGTYRO will help me?
- What are the potential side effects that I may experience?
- How often will I need scans or follow-up visits during treatment?
- How will I know if AUGTYRO is working?
- Will this treatment impact my daily life in any way?

---

---

---

---

---

---

---

---

---

---

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**





Actor portrayals

## HOW TO GET AUGTYRO

You can get AUGTYRO through certain specialty pharmacies that handle treatments, including AUGTYRO, which are not stocked at regular neighborhood pharmacies.

Most specialty pharmacies deliver medications by mail.

Here are some steps your doctor's office and the specialty pharmacy must do before AUGTYRO can be shipped to you:

- Make sure that your health insurance plan has approved AUGTYRO.
- Discuss shipping details with you.
- Collect your co-pay.



Your specialty pharmacy will call to confirm the above details. Make sure you save their number in your phone so you can answer right away or return their calls. This will help you get your AUGTYRO treatment on time.

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**

 **AUGTYRO™**  
(repotrectinib) 40 mg capsules

## At Bristol Myers Squibb, We Provide Support With Purpose

You are the reason behind what we do. The BMS Access Support® program is focused on helping you access your prescribed BMS medicine. If you were prescribed AUGTYRO, you are eligible to enroll in BMS Access Support. The program can help you with:



### Coverage Assistance

Upon your request, BMS Access Support may be able to help you review your insurance benefits and understand your coverage for AUGTYRO.\*

If your insurer requires prior authorization, BMS Access Support may be able to provide you and your doctor information about this requirement.

In the event of a coverage delay or denial, you may be eligible for a bridge program to cover your treatment.† Please call BMS Access Support at 1-800-861-0048 for more information about offerings for your medication.



### Financial Support

If you are an eligible commercially-insured patient, you may pay as little as \$0 per one-month supply.†

If you are insured through a government program or you do not have insurance, BMS Access Support can provide you with information regarding independent charitable foundations.‡



### Free Trial Offer

A Free Trial Offer may be available if you were newly prescribed AUGTYRO.†



### Educational Resources

Understanding the insurance process and accessing your treatment can be challenging. BMS Access Support offers a library of resources that can help you better understand your healthcare coverage and guide you through steps in your treatment journey.

\*The accurate completion and submission of reimbursement- or coverage-related documentation to the patient's insurance plan is the responsibility of the provider and patient. Bristol Myers Squibb and its agents cannot guarantee coverage for any medication or treatment.

†Restrictions apply. Please [click here](#) for full Terms and Conditions, including complete eligibility requirements.

‡It is important to note that charitable foundations are independent from Bristol-Myers Squibb Company and have their own eligibility criteria and evaluation process. Bristol Myers Squibb cannot guarantee that a patient will receive assistance.

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**

#### We're here for you.

Coverage assistance, educational resources, and financial support options may be available through **BMS Access Support**



Call a Patient Access Specialist at **1-800-861-0048**, 8 AM to 8 PM ET, Monday - Friday



Find resources and enrollment information at **[www.BMSAccessSupport.com](http://www.BMSAccessSupport.com)**

**AUGTYRO™**  
(repotrectinib) 40 mg capsules



## Important Facts about AUGTYRO™ (repotrectinib)

This is a summary of important information that you need to know about AUGTYRO. Your healthcare team can work with you to help answer any questions you may have about this medication. [Keep this document in a safe place, so you can refer to it before and during your treatment.](#)

Look out for the following icons as you read:



Talk to your healthcare team



Call a healthcare provider right away



Helpful information to remember

### What is AUGTYRO?

AUGTYRO (repotrectinib) is an oral prescription medicine used to treat **non-small cell lung cancer (NSCLC) in adults** that:

- ✓ Has spread within your chest or to other parts of the body, **AND**
- ✓ Is caused by an **abnormal *ROS1* gene**

It is not known if AUGTYRO is safe and effective in children.



**Talk to your healthcare team** for more information about what it means to have an abnormal *ROS1* gene.

### What is the most important information I should know about AUGTYRO?

AUGTYRO may cause serious side effects. A serious side effect is a side effect that can sometimes become life-threatening and can lead to death. The serious side effects of AUGTYRO may include:

**Central nervous system (CNS) problems.** Symptoms to look out for include:

- dizziness
- vertigo
- changes in mood, such as anxiety, irritability, and depression
- balance or coordination problems
- problems with thinking, such as forgetfulness or confusion
- seeing or hearing things that are not real (hallucinations)
- problems with concentration, attention, memory, and sleep



**Call your healthcare provider right away** if you have any of these symptoms, or any worsening of CNS effects during treatment.

**Lung problems (pneumonitis).** AUGTYRO may cause pneumonitis (inflammation of the lungs), which may be severe or serious. New or worsening symptoms to look out for include:

- Dry cough (without mucus)
- Productive cough (with mucus)
- Wheezing
- Trouble breathing



**Call your healthcare provider right away** if you have any of these symptoms, or any worsening of your current lung problems.

**Liver problems.** Your healthcare provider will do blood tests to check your liver function before starting and during treatment with AUGTYRO. Symptoms of liver problems include:

- your skin or the white part of your eyes turns yellow
- dark or "tea-colored" urine
- light-colored stools (bowel movements)
- loss of appetite
- nausea or vomiting
- pain on the upper right side of your stomach area



**Call your healthcare provider right away** if you develop any of these symptoms of liver problems.

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**

 **AUGTYRO™**  
(repotrectinib) 40 mg capsules

**Muscle problems.** Your healthcare provider will do blood tests before starting treatment with AUGTYRO™, at least every 2 weeks for the first month and as needed during treatment. Symptoms of muscle problems include:

- unexplained muscle pain or muscle pain that does not go away
- tenderness
- weakness



**Call your healthcare provider right away** if you get new or worsening signs and symptoms of muscle problems.

**Increased uric acid level in your blood (hyperuricemia).** AUGTYRO may cause an excess of uric acid in your blood. Your healthcare provider will do tests before and during your treatment with AUGTYRO to check the uric acid level in your blood. Your healthcare provider may prescribe medications if you have high blood uric acid levels. Tell your healthcare provider if you experience symptoms of increased uric acid including:

- red, hot, tender, or swollen joints, especially in your big toe
- decrease in your amount of urine or no urine at all
- nausea or vomiting
- pain in your stomach-area or sides
- pink or brown urine or blood in your urine

**Bone fractures.** AUGTYRO may increase your risk for bone fractures. Bone fractures may happen with or without a fall or other injury.



**Call your healthcare provider right away** if you have pain, changes in movement, or bone abnormalities.

### What should I discuss with my healthcare team before starting treatment?



**Talk to your healthcare provider about all of your medical conditions**, including if you:

- have nervous system (neurological) problems
- have lung or breathing problems other than lung cancer
- have liver problems



**Talk to your healthcare team about all the medicines you are taking**, including:

- prescription medicines
- over-the-counter medicines
- vitamins
- herbal supplements

Certain other medicines may affect how AUGTYRO works, causing side effects.

These are not all the topics you should discuss with your healthcare team. Discuss any questions you may have with your healthcare team.

### What should I discuss with my healthcare team about pregnancy, birth control, and breastfeeding before starting treatment?



**Talk to your healthcare team** if:



**You are pregnant or plan to become pregnant – AUGTYRO may harm your unborn baby**

If you are able to become pregnant, your healthcare team should do a pregnancy test before you start treatment with AUGTYRO.

**If you are able to become pregnant**, you should use effective non-hormonal birth control (contraception) during treatment and for 2 months after your last dose of AUGTYRO.

Birth control methods that contain hormones (such as birth control pills, injections or transdermal system patches) may not work as well during treatment with AUGTYRO.



**Talk to your healthcare team** about **birth control methods** that may be right for you.



**Call your healthcare provider right away** if you become pregnant or think you may be pregnant during treatment with AUGTYRO.

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**

**AUGTYRO™**  
(repotrectinib) 40 mg capsules

### You are a male with a female partner who is able to become pregnant

You should use effective birth control during treatment with AUGTYRO™ and for 4 months after your last dose.



**Call your healthcare provider right away** if your partner becomes pregnant during treatment with AUGTYRO.



### You are breastfeeding or plan to breastfeed

Do not breastfeed during treatment and for 10 days after the last dose of AUGTYRO. It is not known if AUGTYRO passes into your breast milk. Talk to your healthcare team about the best way to feed your baby during this time.

## What are the most common side effects of AUGTYRO?

The most common side effects of AUGTYRO include:

- Dizziness
- Change in sense of taste
- Feeling of numbness or tingling in your arms or legs
- Constipation
- Trouble with balance, coordination, and walking
- Shortness of breath
- Problems with thinking, such as forgetfulness or confusion, memory problems and hallucinations
- Tiredness
- Nausea

These are not all of the possible side effects. **Talk to your doctor** for more information about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA by visiting [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

## How should AUGTYRO be taken?



**Take AUGTYRO exactly as your healthcare provider tells you.** The number of pills that you take will depend on your dose. Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with AUGTYRO if you develop side effects. Do not change your dose or stop taking AUGTYRO without first talking to your healthcare provider.



AUGTYRO is a capsule with a white opaque cap and body, filled with a white to off-white powder. "REP 40" is printed in blue text on the cap.

**Inactive ingredients:** microcrystalline cellulose, sodium lauryl sulfate, croscarmellose sodium, and colloidal silicon dioxide.

**Capsule coating inactive ingredients:** gelatin and titanium dioxide.

**Printing ink:** shellac and FD & C blue #2 aluminum lake.

- ✓ **Do** take AUGTYRO with or without food
- ✓ **Do** take AUGTYRO at about the same time each day, such as in the morning or in the evening\*
- ✓ **Do** swallow AUGTYRO capsules **whole** with water.

*\*Setting an alarm clock or calendar notification is a great way to remember to take medication.*

- ✗ **Do not** open, crush, chew, or dissolve the capsule. **Do not** take broken, cracked, or damaged capsules
- ✗ **Do not** eat grapefruit or drink grapefruit juice during treatment with AUGTYRO
- ✗ **Do not** drive or operate machinery until you know how AUGTYRO affects you and you no longer have symptoms

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**






### What if I miss a dose of AUGTYRO™?

If you miss a dose or vomit at any time after taking a dose of AUGTYRO, **do not take an extra dose.** Just skip the dose and take your next dose at the regularly scheduled time. Do not take 2 doses at the same time to make up a missed or vomited dose.

### How do I store AUGTYRO?

AUGTYRO capsules should be stored at room temperature between 68°F to 77°F (20°C to 25°C). **Keep AUGTYRO and all medicines out of the reach of children.**

 For more information, please click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#) for AUGTYRO. Talk to your healthcare team for more information about this medication.

## MORE NOTES, THOUGHTS, THINGS TO REMEMBER:

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**





Actor portrayals

## SUMMARY OF WARNINGS AND PRECAUTIONS

AUGTYRO (repotrectinib) may cause serious side effects, including central nervous system (CNS) effects, lung problems (pneumonitis), liver problems, muscle problems, increased uric acid level in your blood (hyperuricemia), and bone fractures.

**Please see Important Safety Information on pages 18-21. Click on links for [U.S. Full Prescribing Information](#) and [Patient Information](#).**

 **AUGTYRO™** *with You*  
(repotrectinib) 40 mg capsules

We're here throughout your treatment journey.  
**Scan here to learn more about AUGTYRO**  
and [sign up](#) for AUGTYRO™ *with You* to receive support,  
tools, and educational resources.



**SHARE TO INSPIRE**  
Because your story can help inspire another.

Call us toll-free at **1-855-436-5866**

 Bristol Myers Squibb®

© 2023 Turning Point Therapeutics, Inc., a Bristol-Myers Squibb company.  
AUGTYRO™, AUGTYRO™ *with You*, and the related logos are trademarks of Turning Point Therapeutics, Inc. BMS Access Support® is a registered trademark of Bristol Myers Squibb company. All other trademarks are property of their respective owners.  
3600-US-2300055 12/23

 **AUGTYRO™**  
(repotrectinib) 40 mg capsules